Public reporting burden for the colle maintaining the data needed, and co including suggestions for reducing t VA 22202-4302. Respondents shou does not display a currently valid O	mpleting and reviewing the collection his burden, to Washington Headquald be aware that notwithstanding any	on of information. Send comments reters Services, Directorate for Information	egarding this burden estimate or mation Operations and Reports, 1	any other aspect of thi 1215 Jefferson Davis H	s collection of information, lighway, Suite 1204, Arlington	
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVI 00-00-199	ered 7 to 00-00-1997	
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER					
The Kuroshio/Topo	h Pacific 5b. GRANT NUMBER					
			5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER				
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Florida State University, Department of Oceanography, Tallahassee, FL, 32306-4320				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITOR		10. SPONSOR/MONITOR'S ACRONYM(S)				
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL. Approved for publi		on unlimited				
13. SUPPLEMENTARY NO	ΓES					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	2	PERSON	

Report Documentation Page

Form Approved OMB No. 0704-0188

THE KUROSHIO/TOPOGRAPHY INTERACTION IN THE WESTERN NORTH PACIFIC

Ya Hsueh Department of Oceanography Florida State University Tallahassee, FL 32306-3048

phone: (904)644-670 fax: (904)644-2581 email: hsueh@adin.ocean.fsu.edu Award #: N000149510501

RESEARCH GOALS

The western boundary region of the North Pacific Ocean is unique in that it is porous. Massive exchanges between the Pacific and the marginal seas occur through gaps in the island chain that bounds the open basin. The long term goal of the proposed work is to understand the dynamics of these exchanges and their consequences in terms of ocean variability.

OBJECTIVES

The dynamics of mesoscale flow features that arise from the disruption of the Kuroshio by the continental margin topography in the East China Sea continues to be the focus of the research project. The purpose of the research is to seek an indepth understanding of the dynamics and ramifications of these features.

APPROACH

The examination and interpretation of current-meter, hydrographic, and remote-sensing data collected by Taiwan, Korea, and Japan have continued. In addition, an eddy resolving, general circulation model of the Asian marginal seas is being constructed to address specific dynamic issues of the circulation concerning the Kuroshio/topography interaction. The effort is continued to understand the way the Kuroshio intrudes into and then exit the Luzon Strait.

WORK COMPLETED

To take advantage of considerable current measurements which have begun to become available, an eddy-resolving model of the South China Sea, similar to the expanded Asian marginal seas (AMS) model reported before, has been constructed to study the intrusion of the Kuroshio in the Luzon Strait. Qualitative model/observation comparisons point to an overall agreement and the importance of winds in generating the observed surface current patterns.

RESULTS

- 1. Qualitative comparisons between model outputs and altimeter derived geostrophic velocity normal to a nearly north-south oriented satellite track across the Luzon Strait loop current indicate the model reproduces the overall feature of the loop current, but misses the strong inflow into the South China Sea found at the northern and southern ends of the track. These areas of inflows are obviously associated coastal currents driven by a north wind prevalent during the week (December 29, 1993 January 4, 1994) when the altimetry data were obtained.
- 2. A small leakage of Kuroshio water into the Taiwan Strait is detected in the model. The leakage is found to contribute about half of the total northward volume transport of 0.6 Sy ($1 \text{ Sy} = 10^6 \text{ m}^3/\text{sec}$) found in the model Taiwan Strait.

IMPACT/APPLICATIONS

The qualitative agreement of the South China Sea the geostophic velocity derived from the altimetry data suggests that a working model is in hand for the construction of a model of the Kuroshio intrusion in the Luzon Strait. The foundation is thus in place for a study of the effect of the wind-driven circulation of the South China Sea proper on the exchange of water mass across the Luzon Strait and the northward penetration of Kuroshio water into the Taiwan Strait.

of the effect of the wind-driven circulation of the South China Sea proper on the exchang
of water mass across the Luzon Strait and the northward penetration of Kuroshio water
into the Taiwan Strait.
TRANSITIONS

RELATED PROJECTS

None.

None.

REFERENCES

None.